

PARENTERAL Starter Feeding Regimen for Central Line (Regimen 3a & 3b)

Please refer to dietitian on next working day for prescribed regimen

Is patient at refeeding risk?

- ◆ No - Nutritional risk score <4
Implement Regimen 3a
- ◆ YES - Nutritional risk score \geq 4
Implement Regimen 3b

Aim to continue NG feed 10ml/hr if possible. Reassess regularly for EN eligibility.

Regimen 3a

- ◆ Commence Kabiven 11 @ 72ml per hour over 24hrs.
- ◆ Prescribe Pabrinex IV (one pair) once daily until vitamins & trace elements are added to the bag by pharmacy.
- ◆ Monitor biochemistry daily (U&E, Mg & bone profile) and supplement as required.

Regimen 3b

- ◆ Prescribe Pabrinex IV (one pair) once daily 30 minutes prior to starting feeding on day 1 (Prescribed for first 10 days of feeding).
- ◆ Commence Kabiven 11 @ 23ml per hour over 24hrs.
- ◆ Monitor biochemistry daily (U&E, Mg & bone profile) and supplement as required.

- ◆ If patient unlikely to receive dietetic review within 24hrs on day 2 of feeding and refeeding biochemistry (U&E, Mg & bone profile) within normal range consider \uparrow rate of Kabiven 11 to 45ml per hour over 24hrs.
- ◆ Continue to monitor biochemistry daily (U&E, Mg & bone profile) and supplement as required.

**D
R
A
F
T**